

Sketch of Solutions: Masters Comprehensive Exam in Algebra
Week of January 3-7, 2011

- (1) Let G denote a group containing the distinct elements $\{a, b\}$ and satisfying the following properties:
- Every element $x \in G$ can be written as a product $x = g_1 g_2 \cdots g_n$, for some n , where each g_k , for $1 \leq k \leq n$, is equal to a or b .
 - $(ab)^2 = ab$.

Is G isomorphic to a familiar group, or is there too little information to decide?

SOLUTION: $ab = abab \implies ab = 1$ and $b = a^{-1}$. We have $1 \neq a \neq b \neq 1$. It follows that every element x can be written as a power of a and the group is commutative. The map $k \mapsto a^k$ is a homomorphism of \mathbb{Z} onto G . So G is either infinite cyclic (\mathbb{Z}) or finite cyclic (\mathbb{Z}_n). **NOT ENOUGH INFORMATION.**

- (2) Let G denote the group of rigid motions (i.e., reflections and rotations) of the diamond shape. Answer the following:
- What is this group? **SOLUTION:** There is a horizontal reflection x and a vertical one y . There is one rotation $z = xy$ by 180 degrees. There is the identity. That's it. The only group of order 4 with no element of order 4 is $\mathbb{Z}_2 \times \mathbb{Z}_2$.

- (3) Find as many groups of order 12 as you can. **SOLUTION:** Abelian Case: By fundamental theorem of abelian groups \mathbb{Z}_{12} and $\mathbb{Z}_2 \times \mathbb{Z}_6$ are the only possibilities, since there are unique representations $\mathbb{Z}_{d_1} \times \mathbb{Z}_{d_2}$ with d_1 dividing d_2 . It is not possible to have a third integer d_3 divisible by d_2 .

Nonabelian Case: Certainly the dihedral group \mathbb{D}_{12} and the alternating subgroup \mathbb{A}_4 of \mathbb{S}_4 are two familiar possibilities. Are there others? Sylow theory will help. $\mathbb{D}_{12} = \mathbb{Z}_2 \times \mathbb{S}_3$ has a unique 3-Sylow subgroup ($n_3 = 1$) and 2-Sylow subgroups of the form \mathbb{Z}_2^2 ($n_2 = 3$). The \mathbb{A}_4 has $n_2 = 1$ with the 2-Sylow subgroup $\langle (12)(34), (13)(24), (14)(23) \rangle$. Here $n_3 = 4$: $\langle 123 \rangle$, $\langle 124 \rangle$, $\langle 234 \rangle$, and $\langle 134 \rangle$. (Actually if $n_2 = 1$, only the alternating group is possible.) If $n_2 = 3$ and $n_3 = 4$, there is not enough room in the group. If $n_3 = 1$ and $n_2 = 3$, another group is possible besides \mathbb{D}_{12} . One proceeds by considering that the 4-groups might be cyclic. One of these 4 groups generates the other two through conjugacy by nontrivial elements of the unique 3-Sylow subgroup. This is enough to find generators and relations for this case. $G = \langle a, b \mid a^3 = b^2, a^6 = 1, b^{-1}ab = a^{-1} \rangle$.

- (4) Show that $\mathbb{Z} \times \mathbb{Z}$ is isomorphic to a quotient ring of $R = \left\{ \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} : a, b, d \in \mathbb{Z} \right\}$. **SOLUTION:**

Define the surjective homomorphism $R \rightarrow \mathbb{Z} \times \mathbb{Z}$ by $\left\{ \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \right\} \mapsto (a, d)$. The kernel is $\left\{ \begin{pmatrix} 0 & b \\ 0 & 0 \end{pmatrix} \right\} \cong \mathbb{Z}$.

- (5) The "Diophantine equation" $x^2 + y^2 = 3z^2$ has the solutions $x = 0, y = 0, z = 0$. Show that there are no other integer solutions for x, y, z . Hint: First consider solutions in $\mathbb{Z}/4\mathbb{Z}$. **SOLUTION:** Squares in \mathbb{Z}_4 equal 0 or 1. Assume at least one of x, y, z is nonzero. If each of x, y, z is even, we can divide 4 out of the Diophantine equation, until one or more of the numbers x, y, z is odd. No such solution is possible mod 4, hence no nontrivial integer solutions exist.
- (6) Find the splitting field and Galois group for $x^4 - 2$ over the rationals.

SOLUTIONS: Factor the polynomial $f(x) = x^4 - 2$ to get $(x^2 - \alpha)(x^2 + \alpha)$ with $\alpha^2 = 2$. A splitting field \mathbb{K} must contain α and a real algebraic number β such that $\beta^2 = \alpha$. Then $(i\beta)^2 = -\alpha$. So K contains β and $i\beta$, and therefore i with $i^2 = -1$. The splitting field is $\mathbb{K} = \mathbb{Q}[\beta, i] = \mathbb{Q}[\beta][i] = \mathbb{Q}[i][\beta]$. Let G denote the Galois group of $f(x)$. Since $[\mathbb{K} : \mathbb{Q}] = 8 = |G|$, G is a subgroup of S_4 with 8 elements. The automorphism σ with $\beta \mapsto i\beta$ and fixing i has order 4. The automorphism τ with $i \mapsto -i$ and fixing β has order 2. The dihedral group $D_8 = \langle \sigma, \tau \rangle$ since $(\sigma\tau)^2 = id$.