

## Complex Analysis PhD Exam August 1999

Notation:  $\mathbf{C}$  are the complex numbers and  $\mathbf{N} = \{1, 2, 3, \dots\}$  are the natural numbers.

You will find below two sets of problems. Neither one is supposed to be simpler. The first is however more labour intensive.

**Solve three out of the four problems below:**

1. For each  $n = 0, 1, 2, \dots$ , compute

$$\int_{\Gamma} \frac{\sin^n z}{\ln(1+z)} dz$$

where  $\Gamma$  is the positively oriented circle centered around zero of radius  $1/2$  (and  $\ln$  is the principal branch of the logarithm,  $\ln 1 = 0$ ).

2. Find the complex derivative  $F'(z)$  at  $z = 0$  for

$$F(z) = \int_{2/\pi}^{1/\pi} \frac{\sin(e^z/x)}{x} dx.$$

3. Show that there cannot be a sequence of analytic functions  $f_n$  on  $D := \{|z| < 1\}$  that converge uniformly on  $D$  to  $z^2$  and never vanish in  $D$  (i.e.  $f_n(z) \neq 0$  for all  $z \in D$  and  $n \in \mathbf{N}$ ). (Hint: Use the argument principle.)

4. For what values of  $z \in \mathbf{C}$  is the sum of the series

$$\sum_{n=1}^{\infty} \frac{z^n}{(1-nz)}$$

analytic in  $z$ ?

**Solve three out of the four problems below:**

5. Suppose that  $f$  is entire and  $f'(1/n) = \sin(1/n)$  for all  $n \in \mathbf{N}$ . Argue that  $f(z) + \cos(z)$  is a constant function.

6. Let  $f : \mathbf{C} \rightarrow \mathbf{C}$  be entire and such that  $|f(z)| \leq \sqrt{|z|}$  whenever  $|z| \geq 1$ . Show that  $|f(z)| \leq 1$  for all  $z \in \mathbf{C}$ . (Hint: Show first that  $f$  is constant.)

7. Suppose that  $f : \mathbf{C} \rightarrow \mathbf{C}$  is continuous and that, for all  $z_0, z_1 \in \mathbf{C}$  and any piecewise smooth path  $\Gamma$  from  $z_0$  to  $z_1$ , we have

$$\int_{\Gamma} f(z) dz = e^{z_1^2} - e^{z_0^2}.$$

Prove that  $f$  is analytic. Find the formula for  $f$ .

8. Suppose that  $u$  is a non-constant harmonic function on  $\mathbf{C}$ . Show that  $\{u(z) : |z| < 1\}$  is open. (Hint: Use the maximum principle.)