

Instructions: Closed book. No calculator allowed. Show all work and use correct notation to receive full credit! Write legibly.


1. True or False? Circle ONE answer for each.

True or False: If \mathbf{F} is a vector field, then $\text{div } \mathbf{F}$ is a vector field.

True or False: If \mathbf{F} and \mathbf{G} are vector fields, then $\text{curl}(\mathbf{F} \cdot \mathbf{G}) = \text{curl } \mathbf{F} \cdot \text{curl } \mathbf{G}$.

True or False: If S is a sphere and \mathbf{F} is a constant vector field, then $\iint_S \mathbf{F} \cdot d\mathbf{S} = 0$.

2. Given the vectors $\mathbf{v} = \langle 2, 1, -2 \rangle$, $\mathbf{u} = \langle 3, 2, 1 \rangle$, find the lengths of a and b pictured below:



$$a = \frac{\vec{u} \cdot \vec{v}}{\|\vec{v}\|} = \frac{6 + 2 - 2}{3} = 2$$

$$b = \sqrt{\|\vec{u}\|^2 - a^2} = \sqrt{14 - 4} = \sqrt{10}$$

3. (a) Find an equation for the line through the points $(1, 2, 3)$ and $(-1, 5, 4)$.

$$x(t) = 1 - 2t$$

$$y(t) = 2 + 3t$$

$$z(t) = 3 + t$$

- (b) Find an equation for the plane that is perpendicular to the line in part 3a and passes through the point $(4, 0, 1)$.

$$-2(x-4) + 3(y-0) + 1(z-1) = 0$$

$$\text{or } -2x + 3y + z = -7$$

- (c) At what point do the line in part 3a and the plane in part 3b intersect?

$$-2(1-2t) + 3(2+3t) + (3+t) = -7$$

$$-t = 1 \quad t = -1$$

$$x = 3, y = -1, z = 2 \quad (3, -1, 2)$$

4. Given $f(x, y, z) = \frac{x}{1 + xyz}$:

(a) At the point $(1, 0, 2)$ in what direction does f increase most rapidly?

$$\frac{\partial f}{\partial x} = \frac{(1+xy^2) - x(y^2)}{(1+xy^2)^2} \quad \frac{\partial f}{\partial x}(1, 0, 2) = 1$$

$$\frac{\partial f}{\partial y} = -x^2 (1+xy^2)^{-2} \quad \frac{\partial f}{\partial y}(1, 0, 2) = -2 \quad \nabla f(1, 0, 2) = \langle 1, -2, 0 \rangle$$

$$\frac{\partial f}{\partial z} = -x^2 y (1+xy^2)^{-2} \quad \frac{\partial f}{\partial z}(1, 0, 2) = 0$$

(b) At the point $(1, 0, 2)$ what is the rate of change of f in the direction of $\langle 3, -4, 0 \rangle$?


$$D_{\langle 3, -4, 0 \rangle} f(1, 0, 2) = \frac{\langle 1, -2, 0 \rangle \cdot \langle 3, -4, 0 \rangle}{5} = \frac{11}{5}$$

5. Given $x = r \cos \theta$ and that r and θ depend on t in such a way that when $t = 0$: $r = 2, \theta = \pi/4, \frac{dr}{dt} =$

$3, \frac{d\theta}{dt} = \pi$, find $\frac{dx}{dt}$ at $t = 0$.

$$\begin{aligned} \frac{dx}{dt} &= \frac{\partial x}{\partial r} \frac{dr}{dt} + \frac{\partial x}{\partial \theta} \frac{d\theta}{dt} = (\cos \theta) \frac{dr}{dt} - (r \sin \theta) \frac{d\theta}{dt} \\ &= \left(\frac{3}{2} - \pi\right) \frac{\sqrt{2}}{2} \end{aligned}$$

6. Calculate $\int_0^1 \int_{\sqrt{x}}^1 \sqrt{1+y^3} dy dx$.



$$\begin{aligned} &= \int_0^1 \int_0^{y^2} (1+y^3)^{1/2} dx dy \\ &= \int_0^1 y^2 (1+y^3)^{1/2} dy \\ &= \frac{1}{3} \left(\frac{2}{3}\right) (1+y^3)^{3/2} \Big|_0^1 \\ &= \frac{2}{9} (2^{3/2} - 1) \end{aligned}$$

7. Let \mathcal{W} be the region above the cone $z = \sqrt{x^2 + y^2}$ and below the plane $z = 1$. Calculate $\iiint_{\mathcal{W}} x^2 + y^2 dV$.

$$\begin{aligned} \iiint_{\mathcal{W}} x^2 + y^2 dV &= \int_0^{2\pi} \int_0^1 \int_r^1 r^3 dz dr d\theta \\ &= \int_0^{2\pi} \int_0^1 (1-r)r^3 dr d\theta \\ &= \int_0^{2\pi} \left(\frac{1}{4} r^4 - \frac{1}{5} r^5 \right) \Big|_0^1 d\theta = \frac{1}{20} (2\pi) = \frac{\pi}{10}. \end{aligned}$$

8. Verify that the vector field $\mathbf{F}(x, y, z) = \langle 2xy + z, x^2 + 1, x + 2z \rangle$ is conservative and calculate the work done by \mathbf{F} in moving an object from $(2, -1, 1)$ to $(1, 1, 0)$.

$$\begin{aligned} \nabla \times \mathbf{F} &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ 2x+z & x^2+1 & x+2z \end{vmatrix} = \langle 0, 1-1, 2x-2x \rangle \\ &= \mathbf{0}, \text{ conservative.} \end{aligned}$$

$$f(x, y, z) = x^2 y + xz + y + z^2$$

$$\begin{aligned} \text{work} &= f(1, 1, 0) - f(2, -1, 1) = 2 - (-4 + 2 - 1 + 1) \\ &= 4. \end{aligned}$$

9. Calculate $\oint_C -y^2 dx + xy dy$ where C is the counterclockwise oriented simple closed curve consisting of the piece of the parabola $y = 1 - x^2$ between $(-1, 0)$ and $(1, 0)$ together with the piece of the x -axis between $(-1, 0)$ and $(1, 0)$.

greens theorem

$$\oint_C -y^2 dx + xy dy = \iint_D y + 2y dA = \int_{-1}^1 \int_0^{1-x^2} 3y dy dx$$

$$= \int_{-1}^1 \left. \frac{3}{2} y^2 \right|_{y=0}^{y=1-x^2} dx = \frac{3}{2} \left(\int_{-1}^1 1 - 2x^2 + x^4 dx \right)$$


$$= \frac{3}{2} \left(x - \frac{2}{3} x^3 + \frac{x^5}{5} \right) \Big|_{-1}^1 = \frac{8}{5}$$

10. Find the surface area of the part of surface $z = xy$ that lies within the cylinder $x^2 + y^2 = 1$.

$$SA = \iint_D \sqrt{1 + y^2 + x^2} dA$$

$$= \int_0^{2\pi} \int_0^1 \sqrt{r^2 + 1} r dr d\theta$$

$$= \int_0^{2\pi} \left. \frac{1}{3} (r^2 + 1)^{3/2} \right|_0^1 d\theta = \frac{2\pi}{3} (2\sqrt{2} - 1)$$




11. Let S be the part of the paraboloid $z = 4 - x^2 - y^2$ with $z \geq 0$, oriented with upwards pointing normal vector, and let $\mathbf{F}(x, y, z) = \langle -y, x, z \rangle$. Using Stokes' Theorem, calculate $\iint_S (\text{curl}(\mathbf{F})) \cdot d\mathbf{S}$.

$$\text{Stokes: } \iint_S \text{curl } \vec{F} \cdot d\vec{S} = \oint_C \vec{F} \cdot d\vec{r}$$

$$= \oint_C -y dx + x dy + z dz$$

$$= \oint_C -y dx + x dy$$



$$\text{green's} = \iint_D (1) dA$$

$$= 2 \text{ Area}(D) = 2 \pi (2^2) = 8\pi$$

12. Let $\mathbf{F}(x, y, z) = \langle y, x, z^2 \rangle$ and let S be the closed surface consisting of the cone $z = \sqrt{x^2 + y^2}$, $0 \leq z \leq \sqrt{2}$, and the spherical cap $z = \sqrt{4 - x^2 - y^2}$, $\sqrt{2} \leq z \leq 2$. Using the divergence theorem, calculate the flux, $\iint_S \mathbf{F} \cdot d\mathbf{S}$, of \mathbf{F} outwards across S .

divergence thm.

$$\iint_S \vec{F} \cdot d\vec{S} = \iiint_W \text{div } \vec{F} dV = \iiint_W 2z dV$$

$$\stackrel{\text{spherical}}{=} \int_0^{2\pi} \int_0^{\pi/4} \int_0^2 2 \rho \cos \phi \rho^2 \sin \phi d\rho d\phi d\theta$$

$$= 2 \int_0^{2\pi} \int_0^{\pi/4} \left(\frac{1}{4} \rho^4 \Big|_0^2 \right) \cos \phi \sin \phi d\phi d\theta = \frac{1}{2} (2^4) \left(\frac{-1}{2} \cos^2 \phi \Big|_0^{\pi/4} \right) \cdot 2\pi$$

$$= 4\pi$$

